

GORENSTEIN PROJECTIVE DIMENSION WITH RESPECT TO A SEMIDUALIZING MODULE

DIANA WHITE

Dedicated to the memory of Colleen Kilker

ABSTRACT. We introduce and investigate the notion of G_C -projective modules over (possibly non-noetherian) commutative rings, where C is a semidualizing module. This extends Holm and Jørgensen's notion of C -Gorenstein projective modules to the non-noetherian setting and generalizes projective and Gorenstein projective modules within this setting. We then study the resulting modules of finite G_C -projective dimension, showing in particular that they admit G_C -projective approximations, a generalization of the maximal Cohen-Macaulay approximations of Auslander and Buchweitz. Over a local ring, we provide necessary and sufficient conditions for a G_C -approximation to be minimal.

INTRODUCTION

Over a noetherian ring R , Foxby [9], Golod [10], and Vasconcelos [19] independently initiated the study of semidualizing modules (under different names): a module C is semidualizing if $\mathrm{Hom}_R(C, C) \cong R$ and $\mathrm{Ext}_R^{\geq 1}(C, C) = 0$. Examples include the rank 1 free module and a dualizing (canonical) module, when one exists. Golod [10] used these to define G_C -dimension, a refinement of projective dimension, for finitely generated modules. The G_C -dimension of a finitely generated R -module M is the length of the shortest resolution of M by so-called totally C -reflexive modules; see Definition 4.1. Motivated by Enochs and Jenda's extensions in [7] of Auslander and Bridger's G -dimension [2], Holm and Jørgensen [13] have extended this notion to arbitrary modules over a noetherian ring. The current paper provides a unified and generalized treatment of these concepts, in part by removing the noetherian hypothesis. The tools developed in this paper have been particularly useful for investigating the similarities and differences between certain relative cohomology theories [15, 16] and the stability properties of operators on categories [17].

Section 2 is devoted to the study of the G_C -projective R -modules, which are built from projective and C -projective modules; see Definition 2.1. We show that every module that is either projective or C -projective is G_C -projective in Proposition 2.6. In particular, every R -module admits a G_C -projective resolution. Further properties of the class of G_C -projective modules are contained in the following result; see Theorem 2.8.

2000 *Mathematics Subject Classification.* 13D02, 13D05, 13D07, 13D25, 18G20, 18G25.

Key words and phrases. Gorenstein dimensions, G_C -dimensions, G_C -approximations, proper resolutions, strict resolutions, totally C -reflexives, complete resolutions, complete PC -resolutions, semidualizing modules, Bass classes, C -projectives.

Theorem 1. *The class of G_C -projectives is projectively resolving and closed under direct summands. The class of finitely generated G_C -projective R -modules is closed under summands. The set of G_C -projective R -modules admitting a degreewise finite free resolution is finite projectively resolving.*

Section 2 ends with basic properties of the resulting G_C -projective dimension. In particular, we show that, for an R -module M of G_C -projective dimension $n > 0$, the n th kernel in any G_C -projective resolution is G_C -projective.

Within the class of G_C -projective resolutions, the proper ones exhibit particularly good lifting properties; see 1.5. These are the subject of Section 3. Coupled with Proposition 3.4, the following result shows that every module of finite G_C -projective dimension admits a proper G_C -projective resolution; see Theorem 3.6.

Theorem 2. *If M is an R -module with finite G_C -projective dimension, then M admits a strict G_C -projective resolution, that is, a G_C -resolution of the form*

$$0 \rightarrow C \otimes_R P_n \rightarrow \cdots \rightarrow C \otimes_R P_1 \rightarrow G \rightarrow M \rightarrow 0$$

where G is G_C -projective and $P_1, \dots, P_n$ are projective.

These strict G_C -projective resolutions give rise to G_C -projective approximations, which are similar to the maximal Cohen-Macaulay approximations of Auslander and Buchweitz in [3].

Section 4 is concerned with comparing the G_C -projective and totally C -reflexive properties; see Definition 4.1. The next result is Theorem 4.4, which extends a result of Avramov, Buchweitz, Martsinkovsky, and Reiten [5, (4.2.6)].

Theorem 3. *If M and $\text{Hom}_R(M, C)$ admit degreewise finite projective resolutions, then M is G_C -projective if and only if it is totally C -reflexive.*

The paper closes with several results on minimal proper G_C -projective resolutions of finitely generated modules over noetherian local rings.

1. PRELIMINARIES

Throughout this work R is a commutative ring with unity, $\mathcal{X} = \mathcal{X}(R)$ is a class of unital R -modules, and $\mathcal{X}^f$ is the subclass of finitely generated R -modules in $\mathcal{X}$.

Homological dimensions built from resolutions are fundamental to this work. The prototypes are the projective and injective dimensions.

1.1. An R -complex is a sequence of R -module homomorphisms

$$X = \cdots \xrightarrow{\partial_{n+1}^X} X_n \xrightarrow{\partial_n^X} X_{n-1} \xrightarrow{\partial_{n-1}^X} \cdots$$

such that $\partial_{n-1}^X \partial_n^X = 0$ for each integer n ; the n th homology module of X is $H_n(X) = \text{Ker}(\partial_n^X) / \text{Im}(\partial_{n+1}^X)$. A morphism of complexes $\alpha: X \rightarrow Y$ induces homomorphisms $H_n(\alpha): H_n(X) \rightarrow H_n(Y)$, and α is a *quasiisomorphism* when each $H_n(\alpha)$ is bijective.

The complex X is *bounded* if $X_n = 0$ for $|n| \gg 0$; it is *acyclic* if $X_{-n} = 0 = H_n(X)$ for each $n > 0$. When X is acyclic, the natural morphism $X \rightarrow H_0(X) = M$ is a quasiisomorphism, and X is an $\mathcal{X}$ -projective resolution of M if each X_n is in $\mathcal{X}$; in this event, the exact sequence

$$X^+ = \cdots \xrightarrow{\partial_2^X} X_1 \xrightarrow{\partial_1^X} X_0 \rightarrow M \rightarrow 0$$

is the *augmented $\mathcal{X}$ -projective resolution* of M associated to X . Dually, one defines $\mathcal{X}$ -coresolutions and augmented $\mathcal{X}$ -coresolutions. The *$\mathcal{X}$ -projective dimension* of M is defined as

$$\mathcal{X}\text{-pd}_R(M) = \inf\{\sup\{n \mid X_n \neq 0\} \mid X \text{ is an } \mathcal{X}\text{-projective resolution of } M\}.$$

The nonzero modules in $\mathcal{X}$ are precisely the modules of $\mathcal{X}$ -pd 0.

1.2. The class $\mathcal{X}$ is *projectively resolving* if

- (a) $\mathcal{X}$ contains every projective R -module, and
- (b) for every exact sequence of R -modules $0 \rightarrow M' \rightarrow M \rightarrow M'' \rightarrow 0$ with $M'' \in \mathcal{X}$, one has $M \in \mathcal{X}$ if and only if $M' \in \mathcal{X}$.

The class $\mathcal{X}$ is *finite projectively resolving* if

- (a) $\mathcal{X}$ consists entirely of finitely generated R -modules,
- (b) $\mathcal{X}$ contains every finitely generated projective R -module, and
- (c) for every exact sequence of finitely generated R -modules $0 \rightarrow M' \rightarrow M \rightarrow M'' \rightarrow 0$ with $M'' \in \mathcal{X}$, one has $M \in \mathcal{X}$ if and only if $M' \in \mathcal{X}$.

1.3. Consider an exact sequence of R -modules

$$0 \rightarrow M' \rightarrow M \rightarrow M'' \rightarrow 0.$$

The class $\mathcal{X}$ is *closed under extensions* when $M', M'' \in \mathcal{X}$ implies $M \in \mathcal{X}$, *closed under kernels of epimorphisms* when $M, M'' \in \mathcal{X}$ implies $M' \in \mathcal{X}$ and *closed under cokernels of monomorphisms* when $M', M \in \mathcal{X}$ implies $M'' \in \mathcal{X}$.

1.4. Let M be an R -module. If $X \in \mathcal{X}$ and $\phi: X \rightarrow M$ is a homomorphism, the pair (X, ϕ) is an *$\mathcal{X}$ -precover* of M when, for every homomorphism $\psi: Y \rightarrow M$ where $Y \in \mathcal{X}$, there exists a homomorphism $f: Y \rightarrow X$ such that $\phi f = \psi$. Enochs and Jenda introduced this terminology, which can be found in [8].

1.5. An R -complex Z is *$\mathcal{X}$ -proper* if the complex $\text{Hom}_R(Y, Z)$ is exact for each $Y \in \mathcal{X}$. If $\mathcal{X}$ contains R and Z is $\mathcal{X}$ -proper, then Z is exact.

An $\mathcal{X}$ -resolution X of M is *$\mathcal{X}$ -proper* if the augmented resolution X^+ is $\mathcal{X}$ -proper; by [11, (1.8)] $\mathcal{X}$ -proper resolutions are unique up to homotopy. Accordingly, when M admits an $\mathcal{X}$ -proper resolution X and N is an R -module, the *n th relative homology module* and the *n th relative cohomology module*

$$\text{Tor}_n^{\mathcal{X}}(M, N) = H_n(X \otimes_R N) \quad \text{Ext}_{\mathcal{X}}^n(M, N) = H_{-n} \text{Hom}_R(X, N)$$

are well-defined for each integer n .

1.6. A *degreewise finite projective (respectively, free) resolution* of an R -module M is a projective (respectively, free) resolution P of M such that each P_i is a finitely generated projective (respectively, free). Note that M admits a degreewise finite projective resolution if and only if it admits a degreewise finite free resolution. However, it is possible for a module to admit a bounded degreewise finite projective resolution, but not admit a bounded degreewise finite free resolution. For example, if $R = k_1 \oplus k_2$, where k_1 and k_2 are fields, then $M = k_1 \oplus 0$ is a projective R module, but it does not admit a bounded free resolution.

The next result follows from well-known constructions, but the author is unable to locate an elementary reference.

Lemma 1.7. *The class of R -modules admitting a degreewise finite projective (respectively, free) resolution is closed under summands, extensions, kernels of epimorphisms, and cokernels of monomorphisms.* $\square$

1.8. An R -module C is *semidualizing* if

- (a) C admits a degreewise finite projective resolution,
- (b) The natural homothety map $R \rightarrow \text{Hom}_R(C, C)$ is an isomorphism, and
- (c) $\text{Ext}_R^{\geq 1}(C, C) = 0$.

A free R -module of rank one is semidualizing. If R is noetherian and admits a dualizing module D , then D is a semidualizing.

Note that this definition agrees with the established definition when R is noetherian, in which case condition (a) is equivalent to C being finitely generated. Also, since $\text{Hom}_R(C, C) \cong R$ any homomorphism $\phi: C^n \rightarrow C^m$ can be represented by an $m \times n$ matrix with entries in R .

Finally, note that the hypothesis that C admits a degreewise finite free resolution does not imply that R is noetherian. As one example, take R to be a non-noetherian ring and $C = R$. For an example with $C \neq R$, let $Q \rightarrow R$ be a flat local homomorphism of commutative rings, with Q noetherian and R non-noetherian. If C' is semidualizing over Q with degreewise finite projective resolution F , then $C = C' \otimes_Q R$ is semidualizing over the non-noetherian ring R with degreewise finite projective resolution $F \otimes_Q R$.

1.9. Avramov and Martsinkovsky define a general notion of minimality for complexes in [4, §1]: A complex B is *minimal* if every homotopy equivalence $f: B \rightarrow B$ is an isomorphism. Furthermore, by [4, (1.7)] a complex B is minimal if and only if every morphism $f: B \rightarrow B$ homotopic to the identity map on B is an isomorphism.

1.10. Let M , N , and F be R -modules. The *tensor evaluation* homomorphism

$$\omega_{MNF}: \text{Hom}_R(M, N) \otimes_R F \rightarrow \text{Hom}_R(M, N \otimes_R F)$$

is defined by $\omega_{MNF}(\psi \otimes_R f)(m) = \psi(m) \otimes_R f$. It is straightforward to verify that this is an isomorphism when M is a finitely generated free (or projective) R -module.

Lemma 1.11. *Let F be a flat R -module.*

- (a) *If M admits a degreewise finite projective resolution P , then for $i \geq 0$ there are isomorphisms $\text{Ext}_R^i(M, C \otimes_R F) \cong \text{Ext}_R^i(M, C) \otimes_R F$.*
- (b) *If M admits a degreewise finite projective resolution and $\text{Ext}_R^i(M, C) = 0$ for some $i \geq 0$, then $\text{Ext}_R^i(M, C \otimes_R F) = 0$.*
- (c) *If M admits a degreewise finite projective resolution, F is faithfully flat, and $\text{Ext}_R^i(M, C \otimes_R F) = 0$ for some $i \geq 0$, then $\text{Ext}_R^i(M, C) = 0$.*

Proof. (a) The maps ω_{P_iCF} are isomorphisms by 1.10, hence the desired conclusion follows from the flatness of F and the resulting isomorphism of complexes

$$\text{Hom}_R(P, C \otimes_R F) \cong \text{Hom}_R(P, C) \otimes_R F.$$

(b) and (c) These follow directly from (a). $\square$

1.12. An R -module is *C -projective* if it has the form $C \otimes_R P$ for some projective P . Set $\mathcal{P}_C = \mathcal{P}_C(R) = \{C \otimes_R P \mid P \text{ is projective}\}$. These modules are studied extensively (in the non-commutative setting) in [12]. We state for later use a Lemma that follows readily from [12, (3.6, 5.6, 6.8)].

Lemma 1.13. *Consider an exact sequence of R -modules*

$$(1) \quad 0 \rightarrow M' \rightarrow M \rightarrow M'' \rightarrow 0.$$

When M'' is a (finitely generated) C -projective, M' is a (finitely generated) C -projective if and only if M is a (finitely generated) C -projective. If all of the modules in (1) are C -projective, then (1) splits. $\square$

1.14. The Bass class with respect to C , denoted $\mathcal{B}_C$ or $\mathcal{B}_C(R)$, consists of all R -modules N satisfying

- (a) $\text{Ext}_R^{\geq 1}(C, N) = 0$,
- (b) $\text{Tor}_{\geq 1}^R(C, \text{Hom}_R(C, N)) = 0$, and
- (c) The evaluation map $\nu_{CN}: C \otimes_R \text{Hom}_R(C, N) \rightarrow N$ is an isomorphism.

2. G_C -PROJECTIVE MODULES

In this section we define and develop properties of G_C -projective R -modules and the associated G_C -projective dimension. We begin with a definition which extends the notion of G_C -projective modules found in [13] (where they are referred to as C -Gorenstein projective modules) to the non-noetherian setting.

Definition 2.1. A complete PC -resolution is an exact sequence of R -modules

$$(2) \quad X = \cdots \rightarrow P_1 \rightarrow P_0 \rightarrow C \otimes_R Q^0 \rightarrow C \otimes_R Q^1 \rightarrow \cdots$$

where each P_i and Q^i is projective, and such that the complex $\text{Hom}_R(X, C \otimes_R Q)$ is exact for each projective R -module Q .

An R -module M is G_C -projective if there exists a complete PC -resolution as in (2) with $M \cong \text{Coker}(P_1 \rightarrow P_0)$.

Note that when $C = R$, the definitions above correspond to the definitions of complete resolutions and Gorenstein projective modules. The definition immediately gives rise to the following, which generalizes [11, (2.3)].

Proposition 2.2. *A module M is G_C -projective if and only if $\text{Ext}_R^{\geq 1}(M, C \otimes_R P) = 0$ and M admits a $\mathcal{P}_C$ -coresolution Y with $\text{Hom}_R(Y, C \otimes_R Q)$ exact for any projective Q . $\square$*

Observation 2.3. If M is a G_C -projective R -module, then M admits a complete PC -resolution of the form

$$(3) \quad \cdots \rightarrow F_1 \rightarrow F_0 \rightarrow C \otimes_R F^0 \rightarrow C \otimes_R F^1 \rightarrow \cdots$$

where each F_i and F^i is free. To construct such a sequence from a given complete PC -resolution argue as in [11, (2.4)].

When X is a complex of the form (2), then the complex $\text{Hom}_R(X, C \otimes_R Q)$ is exact for all projective R -modules Q if and only if the complex $\text{Hom}_R(X, C \otimes_R F)$ is exact for all free R -modules F . One implication is immediate. For the other, note that if $Q \oplus Q'$ is free, then we have the following isomorphism of complexes $\text{Hom}_R(X, C \otimes_R (Q \oplus Q')) \cong \text{Hom}_R(X, C \otimes_R Q) \oplus \text{Hom}_R(X, C \otimes_R Q')$.

The next three results provide ways to create G_C -projective modules.

Proposition 2.4. *If X_λ is a collection of complete PC -resolutions, then $\coprod_\lambda X_\lambda$ is a complete PC -resolution. Thus, the class of (finitely generated) G_C -projective R -modules is closed under (finite) direct sums.*

Proof. For any projective R -module Q there is an isomorphism,

$$\mathrm{Hom}_R\left(\coprod_{\lambda} X_{\lambda}, C \otimes_R Q\right) \cong \prod_{\lambda} \mathrm{Hom}_R(X_{\lambda}, C \otimes_R Q).$$

Thus, if the complex $\mathrm{Hom}_R(X_{\lambda}, C \otimes_R Q)$ is exact for all λ then the complex $\mathrm{Hom}_R(\coprod_{\lambda} X_{\lambda}, C \otimes_R Q)$ is exact. It follows that a (finite) direct sum of (finitely generated) G_C -projective R -modules is a (finitely generated) G_C -projective R -module. $\square$

Lemma 2.5. *Let P and Q be projective R -modules and X a complex of R -modules. If the complex $\mathrm{Hom}_R(X, C \otimes_R Q)$ is exact, then the complex $\mathrm{Hom}_R(P \otimes_R X, C \otimes_R Q)$ is exact. Thus, if X is a complete PC -resolution of an R -module M , then $P \otimes_R X$ is a complete PC -resolution of $P \otimes_R M$. The converses hold when P is faithfully projective.*

Proof. Assume the complex $\mathrm{Hom}_R(X, C \otimes_R Q)$ is exact. Since $\mathrm{Hom}_R(P, -)$ is an exact functor, the isomorphism of complexes given by Hom-tensor adjointness

$$\mathrm{Hom}_R(P \otimes_R X, C \otimes_R Q) \cong \mathrm{Hom}_R(P, \mathrm{Hom}_R(X, C \otimes_R Q))$$

implies that $\mathrm{Hom}_R(P \otimes_R X, C \otimes_R Q)$ is exact. It is now straightforward to see that if X is a complete PC -resolution of an R -module M , then $P \otimes_R X$ is a complete PC -resolution of $P \otimes_R M$.

If P is faithfully projective, then the complex $\mathrm{Hom}_R(P, \mathrm{Hom}_R(X, C \otimes_R Q))$ is exact if and only if the complex $\mathrm{Hom}_R(X, C \otimes_R Q)$ is exact. $\square$

Proposition 2.6. *If P is R -projective, then P and $C \otimes_R P$ are G_C -projective. Thus, every R -module admits a G_C -projective resolution.*

Proof. Using Lemma 2.5, it suffices to construct complete PC -resolutions of C and R . By definition, C admits an augmented degree-wise finite free resolution

$$X = \cdots \rightarrow R^{\beta_1} \rightarrow R^{\beta_0} \rightarrow C \rightarrow 0$$

and this is a complete PC -resolution of C . Indeed, the complex X is exact by definition and $C \cong \mathrm{Coker}(R^{\beta_1} \rightarrow R^{\beta_0})$. Furthermore, the complex $\mathrm{Hom}_R(X, C \otimes_R Q)$ is exact for all projective R -modules Q by Lemma 1.11(b), because $\mathrm{Ext}_R^{\geq 1}(C, C) = 0$. Thus, C is G_C -projective.

We now show that

$$\mathrm{Hom}_R(X, C) = 0 \rightarrow R \rightarrow C^{\beta_0} \rightarrow C^{\beta_1} \rightarrow \cdots$$

is a complete PC -resolution of R . First, left exactness of $\mathrm{Hom}_R(-, C)$ and the equality $\mathrm{Ext}_R^{\geq 1}(C, C) = 0$ imply $\mathrm{Hom}_R(X, C)$ is exact. Moreover, since $\mathrm{Hom}_R(X, C)$ consists of finitely presented modules, for any projective R -module Q , tensor evaluation provides the first isomorphism of complexes

$$\mathrm{Hom}_R(\mathrm{Hom}_R(X, C), C \otimes_R Q) \cong \mathrm{Hom}_R(\mathrm{Hom}_R(X, C), C) \otimes_R Q \cong X \otimes_R Q.$$

The second isomorphism follows from the fact that $\mathrm{Hom}_R(C, C) \cong R$. These complexes are exact since the complex X is exact and Q is flat.

Finally, since the class of G_C -projective R -modules contains the class of projective R -modules, every R -module admits a G_C -projective resolution. $\square$

When $C = R$, the following proposition is contained in [11, (2.3)]. The proof is similar to that of [4, (2.2)].

Proposition 2.7. *If X is a complete PC -resolution and L is an R -module admitting a bounded $\mathcal{P}_C$ -projective resolution, then the complex $\text{Hom}(X, L)$ is exact. Thus, if M is G_C -projective, then $\text{Ext}_R^{\geq 1}(M, L) = 0$. $\square$*

The following result is Theorem 1 from the introduction.

Theorem 2.8. *The class of G_C -projectives is projectively resolving and closed under direct summands. The class of finite G_C -projective R -modules is closed under summands. The class of G_C -projective R -modules admitting a degreewise finite projective resolution is finite projectively resolving.*

Proof. Consider an exact sequence

$$(4) \quad 0 \rightarrow M' \xrightarrow{\iota} M \xrightarrow{\rho} M'' \rightarrow 0$$

of R -modules. First, assume that M' and M'' are G_C -projective with complete PC -resolutions X' and X'' , respectively. Use the Horseshoe Lemmas in [11, (1.7)] and [14, (6.20)], together with the fact that the classes of projective and C -projective R -modules are closed under extensions to construct a complex

$$X = \cdots \rightarrow P_1 \rightarrow P_0 \rightarrow C \otimes_R Q^0 \rightarrow C \otimes_R Q^1 \rightarrow \cdots$$

with P_i and Q^i projective and a degreewise split exact sequence of complexes

$$0 \rightarrow X' \rightarrow X \rightarrow X'' \rightarrow 0$$

such that $\text{Coker}(P_1 \rightarrow P_0) \cong M$. To show that M is G_C -projective, it suffices to show that $\text{Hom}_R(X, C \otimes_R Q)$ is exact for all projective R -modules Q . The sequence

$$0 \rightarrow \text{Hom}_R(X'', C \otimes_R Q) \rightarrow \text{Hom}_R(X, C \otimes_R Q) \rightarrow \text{Hom}_R(X', C \otimes_R Q) \rightarrow 0$$

is an exact sequence of complexes. Since the outer two complexes are exact, the associated long exact sequence in homology shows that the middle one is also exact.

Next, assume that M and M'' are G_C -projective with complete PC -resolutions X and X'' , respectively. Comparison lemmas for resolutions, see e.g. [11, (1.8)] and by [14, (6.9)], provide a morphism of chain complexes $\phi: X \rightarrow X''$ inducing ρ on the degree 0 cokernels. By adding complexes of the form $0 \rightarrow P_i'' \xrightarrow{\text{id}} P_i'' \rightarrow 0$ and $0 \rightarrow C \otimes_R (Q^i)'' \xrightarrow{\text{id}} C \otimes_R (Q^i)'' \rightarrow 0$ to X , one can assume ϕ is surjective. Since both the class of projective and C -projective modules are closed under kernels of epimorphisms, see Theorem 1.13, the complex $X' = \ker(\phi)$ has the form

$$X' = \cdots \rightarrow P'_1 \rightarrow P'_0 \rightarrow C \otimes_R (Q^0)' \rightarrow C \otimes_R (Q^1)' \rightarrow \cdots$$

with P'_i and $(Q^i)'$ projective. The exact sequence $0 \rightarrow X' \rightarrow X \rightarrow X'' \rightarrow 0$ is degreewise split by Lemma 1.13, so an argument similar to that of the previous paragraph implies that X' is a complete PC -resolution and M' is G_C -projective.

Since the class of G_C -projective R -modules is projectively resolving by the previous paragraphs and closed under arbitrary direct sums by Proposition 2.4, it follows from Eilenberg's swindle [11, (1.4)] that they are also closed under direct summands.

When the exact sequence (4) consists of modules admitting a degreewise finite projective resolution, one can check that the above constructions can be carried out using finite modules. Finally, if G is a finitely generated G_C -projective, then any summand is also G_C -projective. Since summands of finitely generated modules are finitely generated, this implies that the class of finitely generated G_C -projective modules is closed under summands. $\square$

When $C = R$, the next proposition follows readily from the symmetry of the definition of the Gorenstein projectives. However, the case of G_C -projectives, the situation is more subtle. Nonetheless, significant symmetry exists.

Proposition 2.9. *Every cokernel in a complete PC -resolution is G_C -projective.*

Proof. Consider a complete PC -resolution

$$(5) \quad X = \cdots \rightarrow P_1 \rightarrow P_0 \rightarrow C \otimes_R Q^0 \rightarrow C \otimes_R Q^1 \rightarrow \cdots$$

and set $M = \text{Coker}(P_1 \rightarrow P_0)$ and $K = \text{Coker}(P_2 \rightarrow P_1)$. Since M and P_0 are G_C -projective, the exact sequence

$$0 \rightarrow K \rightarrow P_0 \rightarrow M \rightarrow 0$$

shows that K is G_C -projective; see Theorem 2.8. Inductively, one can show that $\text{Coker}(P_{i+1} \rightarrow P_i)$ is G_C -projective for every positive integer i .

Set $N_{-1} = M$, $N_0 = \text{Coker}(P_0 \rightarrow C \otimes_R Q^0)$, and $N_i = \text{Coker}(C \otimes_R Q^{i-1} \rightarrow C \otimes_R Q^i)$ for $i \geq 1$. Using Proposition 2.2, we will be done once we verify that $\text{Ext}_R^{\geq 1}(N_i, C \otimes_R Q) = 0$ for all projective R -modules Q . For each $i > -1$, consider the exact sequence

$$Y_i = 0 \rightarrow N_i \rightarrow C \otimes_R Q^{i+1} \rightarrow N_{i+1} \rightarrow 0.$$

By induction, one has $\text{Ext}_R^{\geq 1}(N_i, C \otimes_R Q) = 0$. Proposition 2.6 implies that $C \otimes_R Q^{i+1}$ is G_C -projective for each $i \geq 0$, and hence $\text{Ext}_R^{\geq 1}(C \otimes_R Q^{i+1}, C \otimes_R Q) = 0$. The long exact sequence in $\text{Ext}_R(-, C \otimes_R Q)$ associated to Y_i provides $\text{Ext}_R^{\geq 2}(N_{i+1}, C \otimes_R Q) = 0$. Furthermore, since $\text{Hom}_R(X, C \otimes_R Q)$ is exact, so is the complex $\text{Hom}_R(Y_i, C \otimes_R Q)$. Therefore, since $\text{Ext}_R^1(C \otimes_R Q^{i+1}, C \otimes_R Q) = 0$, one has $\text{Ext}_R^1(N_{i+1}, C \otimes_R Q) = 0$. $\square$

The class of G_C -projective R -modules can be used to define the G_C -projective dimension, denoted $\text{G}_C\text{-pd}_R(-)$; see 1.1. The following 5 results are proved similarly to [11, (2.18), (2.19), (2.20), (2.21), (2.24)]. We collect them here for ease of reference.

Proposition 2.10. *Let $0 \rightarrow K \rightarrow G \rightarrow M \rightarrow 0$ be an exact sequence of R -modules where G is G_C -projective. If M is G_C -projective, then so is K . Otherwise, one has $\text{G}_C\text{-pd}_R(K) = \text{G}_C\text{-pd}_R(M) - 1$.* $\square$

Proposition 2.11. *If $(M_\lambda)_{\lambda \in \Lambda}$ is a collection of R -modules, then*

$$\text{G}_C\text{-pd}_R\left(\coprod_{\lambda} M_\lambda\right) = \sup\{\text{G}_C\text{-pd}_R(M_\lambda) \mid \lambda \in \Lambda\}. \quad \square$$

Proposition 2.12. *Let M be an R -module such that $\text{G}_C\text{-pd}_R(M)$ is finite and let n be an integer. The following are equivalent.*

- (i) $\text{G}_C\text{-pd}_R(M) \leq n$.
- (ii) $\text{Ext}_R^i(M, L) = 0$ for all $i > n$ and all R -modules L with $\mathcal{P}_C\text{-pd}(L) < \infty$.
- (iii) $\text{Ext}_R^i(M, C \otimes_R P) = 0$ for all $i > n$ and all projective R -modules P .
- (iv) In every exact sequence $0 \rightarrow K_n \rightarrow G_{n-1} \rightarrow \cdots \rightarrow G_0 \rightarrow M \rightarrow 0$ where the G_i are G_C -projective, one has that K_n is also G_C -projective. $\square$

Proposition 2.13. *Let M be an R -module with $\text{G}_C\text{-pd}_R(M) < \infty$. If M admits a degreewise finite projective resolution, then there is an equality $\text{G}_C\text{-pd}_R(M) = \sup\{i \in \mathbb{Z} \mid \text{Ext}_R^i(M, C) \neq 0\}$.* $\square$

Proposition 2.14. *If two modules in an exact sequence have finite G_C -projective dimension, then so does the third.* $\square$

When $C = R$, there are numerous proofs (see e.g. [4, (3.4)] or [11, (2.27)]) of the following: if M is an R -module of finite projective dimension, then there is an equality $\text{pd}_R(M) = \text{G-pd}_R(M)$. Since G_C -dimension can be viewed as a refinement of projective dimension, it makes sense to ask the following:

Question 2.15. If M is an R -module of finite projective dimension, must $\text{pd}_R(M) = \text{G}_C\text{-pd}_R(M)$?

Over a noetherian, local ring, the affirmative answer in the case of finitely generated modules follows immediately from the AB-formulas for projective dimension and G_C -dimension. Over a non-local noetherian ring, an affirmative answer follows from work in [11] and [13]. However, as of the writing of this paper, the author does not know the answer to this question in general.

However, arguably the more natural comparison is between $\mathcal{P}_C$ -dimension and G_C -dimension. We have the following.

Proposition 2.16. *If M is an R -module of finite $\mathcal{P}_C$ -projective dimension, then $\mathcal{P}_C\text{-pd}_R(M) = \text{G}_C\text{-pd}_R(M)$.*

Proof. Using Proposition 2.12, it suffices to show that if M is G_C -projective with finite $\mathcal{P}_C$ -projective dimension, then M is C -projective. To this end, consider an exact sequence of the form

$$0 \rightarrow K \rightarrow C \otimes_R P \rightarrow M \rightarrow 0$$

where P is projective and $\text{G}_C\text{-pd}_R(K) < \infty$. By Proposition 2.12, $\text{Ext}_R^1(M, K) = 0$ so the above sequence splits, forcing M to be a summand of $C \otimes_R P$. Since the class of C -projectives is closed under summands by 1.13, this implies that M is C -projective, as desired. $\square$

3. G_C -PROJECTIVE RESOLUTIONS AND APPROXIMATIONS

In this section we prove the existence of strict and proper G_C -projective resolutions and of G_C -projective approximations. These will give rise to well-defined relative (co)homology functors, see Remark 3.7, which are further studied in [15] and [16]. We begin with the requisite definitions.

Definition 3.1. Let M be an R -module of finite G_C -projective dimension. A *strict G_C -projective resolution* of M is a bounded G_C -projective resolution G such that for $i \geq 1$, there exists a projective R -module P_i such that $G_i \cong C \otimes_R P_i$. This gives rise to an associated *G_C -projective approximation* of M ; that is, an exact sequence of R -modules

$$0 \rightarrow K \rightarrow G \rightarrow M \rightarrow 0$$

in which $\mathcal{P}_C\text{-dim}_R(K)$ is finite and G is G_C -projective.

We provide two examples. The first corresponds to the situation when C is dualizing, the second to when $C = R$.

Example 3.2. When R is a local Cohen-Macaulay ring with dualizing module D , Auslander and Buchweitz [3] show that every finitely generated module M admits a maximal Cohen-Macaulay approximation, that is, an exact sequence of the form

$$0 \rightarrow K \rightarrow G \rightarrow M \rightarrow 0$$

where K has finite injective dimension and G is maximal Cohen-Macaulay. This gives rise to a resolution of the form

$$0 \rightarrow D^{\alpha_n} \rightarrow \cdots \rightarrow D^{\alpha_0} \rightarrow G \rightarrow M \rightarrow 0$$

where G is a maximal Cohen-Macaulay module.

Example 3.3. When R is noetherian and M is an R -module of finite G -dimension, Avramov and Martsinkovsky [4, (3.8)] and Holm [11, (2.10)] provide several constructions of G -approximations, that is, exact sequences of the form

$$0 \rightarrow K \rightarrow G \rightarrow M \rightarrow 0$$

where K has finite projective dimension and G is totally reflexive (see 4.1). These give rise to strict G -approximations, namely, exact sequences of the form

$$0 \rightarrow R^{\alpha_n} \rightarrow \cdots \rightarrow R^{\alpha_0} \rightarrow G \rightarrow M \rightarrow 0$$

where G is totally reflexive.

The existence of strict G_C -projective resolutions implies the existence of proper G_C -projective resolutions.

Proposition 3.4. *Augmented strict G_C -projective resolutions are G_C -proper.*

Proof. Let H be a G_C -projective R -module and

$$(6) \quad 0 \rightarrow C \otimes_R P_n \rightarrow \cdots \rightarrow C \otimes_R P_1 \rightarrow G \rightarrow M \rightarrow 0$$

an augmented strict G_C -projective resolution. Since $\text{Ext}_R^1(H, C \otimes_R P_n) = 0$ by Proposition 2.12, applying $\text{Hom}_R(H, -)$ to the exact sequence $0 \rightarrow C \otimes_R P_n \rightarrow C \otimes_R P_{n-1} \rightarrow K_{n-2} \rightarrow 0$ provides an exact sequence

$$0 \rightarrow \text{Hom}_R(H, C \otimes_R P_n) \rightarrow \text{Hom}_R(H, C \otimes_R P_{n-1}) \rightarrow \text{Hom}_R(H, K_{n-2}) \rightarrow 0.$$

Continuing to break the exact sequence (6) into short exact sequences and applying Proposition 2.12 shows that (6) is G_C -proper. $\square$

The existence of a strict G_C -projective resolution for a module M of finite G_C -projective dimension which is in the Bass class of R with respect to C (see 1.14) was shown in [13, (5.9)]. We offer an alternative construction, motivated by [3], that has the added advantage of not requiring any Bass class assumption. When R is noetherian and M is finitely generated, this is [1, (2.13)]. We begin by proving a lemma.

Lemma 3.5. *Let $\phi: G \rightarrow V$ be a homomorphism between G_C -projective R -modules. If $0 \rightarrow G \xrightarrow{\psi} U \rightarrow N \rightarrow 0$ is an exact sequence of R -modules such that N is G_C -projective then the pushout module H of the maps ϕ and ψ is G_C -projective.*

Proof. We have a commutative diagram with exact rows

$$\begin{array}{ccccccc} 0 & \longrightarrow & G & \xrightarrow{\psi} & U & \longrightarrow & N \longrightarrow 0 \\ & & \downarrow \phi & & \downarrow & & \downarrow = \\ 0 & \longrightarrow & V & \longrightarrow & H & \longrightarrow & N \longrightarrow 0. \end{array}$$

Since N and V are G_C -projective, Proposition 2.8 implies H is G_C -projective. $\square$

The next result contains Theorem 2 from the introduction.

Theorem 3.6. *If M is an R -module with finite G_C -projective dimension, then M admits a strict G_C -projective resolution and hence a G_C -projective approximation.*

Proof. Assume $G_C\text{-pd}_R(M) = n$. By Proposition 2.12, truncating an augmented free resolution of M yields an augmented G_C -projective resolution of M

$$0 \rightarrow G_n \xrightarrow{\phi_n} F_{n-1} \rightarrow \cdots \rightarrow F_0 \rightarrow M \rightarrow 0$$

A complete PC -resolution of G_n gives rise to an exact sequence

$$0 \rightarrow G_n \xrightarrow{\psi} C \otimes_R P_n \rightarrow N \rightarrow 0$$

where P_n is projective and N is G_C -projective. Lemma 3.5 provides a commutative diagram (note that the orientation is not the same as in the previous lemma)

$$\begin{array}{ccccccc} 0 & \longrightarrow & G_n & \xrightarrow{\phi_n} & F_{n-1} & \longrightarrow & F_{n-2} \longrightarrow \cdots \\ & & \downarrow \psi & & \downarrow & & \\ 0 & \longrightarrow & C \otimes_R P_n & \xrightarrow{\phi'} & G_{n-1} & & \end{array}$$

with exact rows in which G_{n-1} is G_C -projective. As G_{n-1} is a pushout module, the maps ϕ_n and ϕ' have isomorphic cokernels, resulting in a G_C -resolution

$$0 \rightarrow C \otimes_R P_n \xrightarrow{\phi'} G_{n-1} \rightarrow F_{n-2} \rightarrow \cdots \rightarrow F_0 \rightarrow 0.$$

Continuing this process yields a strict G_C -projective resolution of M . $\square$

Remark 3.7. As noted in the introduction, Proposition 3.4 and Theorem 3.6 imply the following: every module M of finite G_C -projective dimension admits a proper G_C -projective resolution. Hence, the relative (co)homology functors $\text{Ext}_{G_C}^n(M, -)$ and $\text{Tor}_n^{G_C}(M, -)$ are well-defined for each integer n ; see 1.5.

We close the section with a complement to Proposition 2.11, which is proved as in [11, (2.11)].

Corollary 3.8. *Let $0 \rightarrow G' \rightarrow G \rightarrow M \rightarrow 0$ be an exact sequence of R -modules. Assume G and G' are G_C -projective and that $\text{Ext}_R^1(M, C \otimes_R Q) = 0$ for all projective R -modules Q . Then M is G_C -projective.* $\square$

4. CONNECTIONS WITH TOTALLY C -REFLEXIVE MODULES.

In this section, we reconnect with Golod's G_C -dimension.

Definition 4.1. Let M be an R -module and assume that M and $\text{Hom}_R(M, C)$ admit a degreewise finite projective resolution. The module M is *totally C -reflexive* if the following conditions hold

- (a) The natural biduality map $M \rightarrow \text{Hom}_R(\text{Hom}_R(M, C), C)$ is an isomorphism,
- (b) $\text{Ext}_R^{\geq 1}(M, C) = 0$, and
- (c) $\text{Ext}_R^{\geq 1}(\text{Hom}_R(M, C), C) = 0$.

Observation 4.2. Finitely generated free modules are totally C -reflexive, as is the R -module C^n for any positive integer n . If M is totally C -reflexive, then it is straightforward to check that any summand M' of M is also totally C -reflexive (using Lemma 1.7 to see that M' admits a degreewise finite free resolution). Thus, finitely generated projective R -modules are also totally C -reflexive, and so every finitely generated R -module admits a resolution by totally C -reflexive modules.

When R is noetherian, the homological dimension which arises by resolving a given module by totally C -reflexive modules is known as the G_C -dimension of a module, which was first introduced by Golod; see [10]. In the case $C = R$, this is Auslander and Bridger's G -dimension [2].

Next we provide a useful characterization of totally C -reflexive modules, which generalizes [5, (4.1.4)].

Lemma 4.3. *An R -module M is totally C -reflexive if and only if there is an exact sequence of the form*

$$(7) \quad X = \cdots \rightarrow R^{\beta_1} \rightarrow R^{\beta_0} \rightarrow C^{\alpha_0} \rightarrow C^{\alpha_1} \rightarrow \cdots$$

with $M \cong \text{Coker}(R^{\beta_1} \rightarrow R^{\beta_0})$ and such that $\text{Hom}_R(X, C)$ is exact.

Proof. Set $(-)^{\dagger} = \text{Hom}_R(-, C)$. Assume first that M is totally C -reflexive. By definition, there exists augmented degreewise finite free resolutions

$$F = \cdots \rightarrow R^{\beta_1} \rightarrow R^{\beta_0} \rightarrow M \rightarrow 0$$

$$G = \cdots \rightarrow R^{\alpha_1} \rightarrow R^{\alpha_0} \rightarrow M^{\dagger} \rightarrow 0.$$

The complexes $F^{\dagger}$ and $G^{\dagger}$ are exact, as $\text{Ext}_R^{\geq 1}(M, C) = 0 = \text{Ext}_R^{\geq 1}(M^{\dagger}, C)$. The isomorphism $M \cong M^{\dagger\dagger}$ shows that $G^{\dagger}$ has the form

$$G^{\dagger} \cong 0 \rightarrow M \rightarrow C^{\alpha_0} \rightarrow C^{\alpha_1} \rightarrow \cdots.$$

Splicing together the complexes F and $G^{\dagger}$ provides an exact sequence X of the form (7) with $M \cong \text{Coker}(R^{\beta_1} \rightarrow R^{\beta_0})$. The fact that $F^{\dagger}$ and G are exact implies that $X^{\dagger}$ is exact.

Conversely, assume that M admits a resolution X of the form (7) such that

$$(8) \quad X^{\dagger} = \cdots \rightarrow R^{\alpha_1} \rightarrow R^{\alpha_0} \rightarrow C^{\beta_0} \rightarrow C^{\beta_1} \rightarrow \cdots$$

is exact. Consider the following “soft truncations” of X

$$F = \cdots \rightarrow R^{\beta_1} \rightarrow R^{\beta_0} \rightarrow M \rightarrow 0$$

$$H = 0 \rightarrow M \rightarrow C^{\alpha_0} \rightarrow C^{\alpha_1} \rightarrow \cdots.$$

The complex $X^{\dagger}$ is exact and therefore so are $F^{\dagger}$ and $H^{\dagger}$.

Since F is an augmented free resolution of M , this implies that $\text{Ext}_R^{\geq 1}(M, C) = 0$. The biduality maps and exactness of $H^{\dagger}$ provide a commutative diagram

$$\begin{array}{ccccccc} H = 0 & \longrightarrow & M & \longrightarrow & C^{\alpha_0} & \longrightarrow & C^{\alpha_1} \longrightarrow \cdots \\ & & \downarrow \delta_M^C & & \downarrow \delta_{C^{\alpha_0}}^C & & \downarrow \delta_{C^{\alpha_1}}^C \\ H^{\dagger\dagger} = 0 & \longrightarrow & \text{Hom}_R(\text{Hom}_R(M, C), C) & \longrightarrow & C^{\alpha_0} & \longrightarrow & C^{\alpha_1} \longrightarrow \cdots \end{array}$$

The top row is exact by definition, while a routine diagram chase and the fact that $\text{Hom}_R(-, C)$ is left exact shows that the bottom row is exact. Since $\delta_{C^{\alpha_1}}^C$ and $\delta_{C^{\alpha_0}}^C$ are isomorphisms, the snake lemma implies that the map δ_M^C is an isomorphism. Finally, the exact sequence $H^{\dagger}$ is an augmented degreewise finite free resolution of $M^{\dagger}$. Thus, exactness of $H^{\dagger\dagger}$ implies that $\text{Ext}_R^{\geq 1}(M^{\dagger}, C) = 0$ and thus M is totally C -reflexive. $\square$

The next result is Theorem 3 from the introduction.

Theorem 4.4. *If M and $\text{Hom}_R(M, C)$ admit degreewise finite projective resolutions, then M is G_C -projective if and only if it is totally C -reflexive.*

Proof. Set $(-)^{\dagger} = \text{Hom}_R(-, C)$ and let F and G be degreewise finite free resolutions of M and $\text{Hom}_R(M, C)$, respectively.

Assume first that M is totally C -reflexive. By Lemma 4.3, there is an exact sequence

$$X = \cdots \rightarrow R^{\beta_1} \rightarrow R^{\beta_0} \rightarrow C^{\alpha_0} \rightarrow C^{\alpha_1} \rightarrow \cdots$$

with $M \cong \text{Coker}(R^{\beta_1} \rightarrow R^{\beta_0})$ and such that $\text{Hom}_R(X, C)$ is exact. An argument similar to the one used in the proof of Lemma 1.11 implies that the complex $\text{Hom}_R(X, C \otimes_R P)$ is exact, and so X is a complete PC -resolution of M .

Conversely, assume that M is G_C -projective and let

$$\cdots \rightarrow P_1 \rightarrow P_0 \rightarrow C \otimes_R F^0 \rightarrow C \otimes_R F^1 \rightarrow \cdots$$

be a complete PC -resolution of M in which each F^i is a free R -module; see Observation 2.3. We show M is totally C -reflexive by constructing a complex X as in Lemma 4.3. To this end, it suffices to construct an augmented $\mathcal{P}_C^f$ -coresolution

$$Y = 0 \rightarrow M \rightarrow C \otimes_R R^{\alpha_0} \rightarrow C \otimes_R R^{\alpha_1} \rightarrow \cdots$$

where each α_i is a non-negative integer and $Y^{\dagger}$ is exact. Indeed, Proposition 2.12 implies that $\text{Ext}_R^{\geq 1}(M, C \otimes_R P) = 0$ for any projective R -module P ; in particular $\text{Ext}_R^{\geq 1}(M, C) = 0$. It follows that $(F^+)^{\dagger}$ is exact. Splicing together the complexes F and Y provides the desired complex X .

We now build the complex Y piece by piece. Consider the exact sequence

$$0 \rightarrow M \rightarrow C \otimes_R F^0 \rightarrow G \rightarrow 0$$

arising from the given complete PC -resolution of M . By Proposition 2.9 we know that G is G_C -projective. Since $C \otimes_R F^0$ is a direct sum of copies of C we know that the image of the finitely generated module M is contained in a finite direct sum of copies C . That is, the image of M is contained in a finitely generated submodule $C \otimes_R R^{\alpha_0}$ of $C \otimes_R F^0$. Thus, we have a commutative diagram with exact rows

$$(9) \quad \begin{array}{ccccccc} 0 & \longrightarrow & M & \longrightarrow & C \otimes_R R^{\alpha_0} & \longrightarrow & H \longrightarrow 0 \\ & & \downarrow = & & \downarrow & & \downarrow \\ 0 & \longrightarrow & M & \longrightarrow & C \otimes_R F^0 & \longrightarrow & G \longrightarrow 0. \end{array}$$

Let P be a projective R -module and set $\mathcal{F} = \text{Hom}_R(-, C \otimes_R P)$. Since $C \otimes_R R^{\alpha_0}$ and G are G_C -projective, we have $\text{Ext}_R^1(G, C \otimes_R P) = 0 = \text{Ext}_R^1(C \otimes_R R^{\alpha_0}, C \otimes_R P)$. Hence, applying $\mathcal{F}$ to (9) yields a commutative diagram with exact rows

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathcal{F}(G) & \longrightarrow & \mathcal{F}(C \otimes_R F^0) & \longrightarrow & \mathcal{F}(M) \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow = \\ 0 & \longrightarrow & \mathcal{F}(H) & \longrightarrow & \mathcal{F}(C \otimes_R R^{\alpha_0}) & \longrightarrow & \mathcal{F}(M) \longrightarrow \text{Ext}_R^1(H, C \otimes_R P) \longrightarrow 0. \end{array}$$

A routine diagram chase shows that $\text{Ext}_R^1(H, C \otimes_R P) = 0$. Proposition 2.12 and Proposition 2.14 then imply that H is G_C -projective. Since M and R^{α_0} admit degreewise finite projective resolutions, so does H . Applying $\text{Hom}_R(-, C)$ to the exact sequence

$$0 \rightarrow M \rightarrow C \otimes_R R^{\alpha_0} \rightarrow H \rightarrow 0$$

gives rise to an exact sequence

$$0 \rightarrow \operatorname{Hom}_R(H, C) \rightarrow R^{\alpha_0} \rightarrow \operatorname{Hom}_R(M, C) \rightarrow 0.$$

Here we used the facts that $\operatorname{Hom}_R(C \otimes_R R^{\alpha_0}, C) \cong R^{\alpha_0}$ and $\operatorname{Ext}_R^1(H, C) = 0$ because H is G_C -projective. Since $\operatorname{Hom}_R(M, C)$ and R^{α_0} admit degreewise finite projective resolutions, so does $\operatorname{Hom}_R(H, C)$; see Lemma 1.7. Thus, we can proceed inductively to construct the complex Y with the given properties. $\square$

Corollary 4.5. *If M and $\operatorname{Hom}_R(M, C)$ admit degreewise finite projective resolutions, then M has finite G_C -projective dimension if and only if it has finite G_C -dimension. Moreover, these values coincide.* $\square$

Combining this with the AB-formula for G_C -dimension, see [6, (3.14)], and Proposition 2.16, we have an AB-formula for modules of finite $\mathcal{P}_C$ -dimension.

Corollary 4.6. *Let R be a local, noetherian ring. If M is a finitely generated R -module of finite $\mathcal{P}_C$ -dimension, then $\mathcal{P}_C\text{-pd}_R(M) = \operatorname{depth}(R) - \operatorname{depth}_R(M)$.* $\square$

The next result compares with Theorem 3.6.

Corollary 4.7. *If M and $\operatorname{Hom}_R(M, C)$ admit degreewise finite projective resolutions and $G_C\text{-pd}_R(M)$ is finite, then M admits a strict G_C^f -resolution.* $\square$

We conclude the paper with results on minimal proper G_C -projective resolutions; see 1.9 for the definition of a minimal complex. Note that Proposition 4.10(b) shows, in particular, that such resolutions are strict. We begin with two lemmas, the first of which follows as in [4, (8.1)].

Lemma 4.8. *Over a local, ring R , a complex H consisting of modules in $\mathcal{P}_C^f$ is minimal if and only if $\partial(H) \subseteq \mathfrak{m}H$.* $\square$

Lemma 4.9. *Let R be local, noetherian and M a finitely generated R -module which admits a bounded $\mathcal{P}_C^f$ -resolution. Then M admits a minimal $\mathcal{P}_C^f$ -resolution.*

Proof. An augmented bounded $\mathcal{P}_C^f$ -resolution of M

$$X^+ = 0 \rightarrow C^{\alpha_n} \rightarrow \dots \rightarrow C^{\alpha_1} \rightarrow C^{\alpha_0} \rightarrow M \rightarrow 0.$$

is also an augmented strict G_C -projective resolution of M and so Proposition 3.4 implies that it is proper. Applying the functor $\operatorname{Hom}_R(C, -)$ to X and using the fact that $\operatorname{Hom}_R(C, C) \cong R$ yields an exact sequence

$$\operatorname{Hom}_R(C, X^+) = 0 \rightarrow R^{\alpha_n} \rightarrow \dots \rightarrow R^{\alpha_1} \rightarrow R^{\alpha_0} \rightarrow \operatorname{Hom}_R(C, M) \rightarrow 0.$$

which is an augmented finite free resolution of $\operatorname{Hom}_R(C, M)$. There is an isomorphism of complexes $\operatorname{Hom}_R(C, X) \cong F \oplus G$ where F is an augmented minimal free resolution of $\operatorname{Hom}_R(C, M)$ and G is a contractible complex of free modules. Recall that G is *contractible* if the identity map on G is homotopic to the zero map.

Since M has finite $\mathcal{P}_C$ -dimension, [18, (2.9)] implies that $M \in \mathcal{B}_C(R)$ (see 1.14 for the definition). This provides the first isomorphism below

$$\begin{aligned} X^+ &\cong C \otimes_R \operatorname{Hom}_R(C, X^+) \\ &\cong (C \otimes_R F) \oplus (C \otimes_R G) \end{aligned}$$

while the second follows from the isomorphism $\operatorname{Hom}_R(C, X^+) \cong F \oplus G$ and the fact that finite direct sums commute with tensor products. It is now straightforward to

verify that the complex $C \otimes_R F$ is contractible and that the complex $C \otimes_R F$ is a minimal $\mathcal{P}_C$ -resolution of M , as desired. $\square$

The following structure result is the key to demonstrating the differences between the relative cohomology theories $\text{Ext}_{\mathcal{P}_C}$, Ext_{G_C} , and Ext_R in [15].

Proposition 4.10. *Assume that R is local, and let M be a finitely generated R -module of finite G_C -projective dimension. If M and $\text{Hom}_R(M, C)$ admit degreewise finite projective resolutions, then the following hold.*

- (a) *The module M admits a minimal proper G_C -projective resolution.*
- (b) *A given G_C -projective resolution H of M is minimal if and only if the following conditions hold.*
 - (1) $H_n \cong C^{\alpha_n}$ for all $n \geq 1$,
 - (2) $\partial_n^H(H_n) \subseteq \mathfrak{m}H_{n-1}$ for all $n \geq 2$, and
 - (3) $\partial_1^H(H_1)$ contains no nonzero C -summand of H_0 .

Proof. We begin by showing that a G_C -projective resolution H satisfying conditions (1)–(3) is minimal. First, observe that H_0 is finitely generated because M and H_1 are so. Let $\gamma: H \rightarrow H$ be a morphism that is homotopic to id_H . Using 1.9, we need to show that γ_n is an isomorphism for each integer n .

For $n \geq 0$, let $\theta_n: H_n \rightarrow H_{n+1}$ be maps such that $\gamma_n - \text{id}_{H_n} = \theta_{n-1}\partial_n^H + \partial_{n+1}^H\theta_n$, which exist since γ is homotopic to id_H . For $n \geq 2$, condition (2) implies $\partial_n^H \otimes_R k = 0$, and so $\gamma_n \otimes_R k - \text{id}_{H_n} \otimes_R k = 0$. Nakayama's Lemma implies γ_n is a surjective endomorphism, and hence bijective.

Now let $n = 1$. We verify the containment $\text{Im}(\theta_0\partial_1^H) \subseteq \mathfrak{m}C^{\alpha_1}$ and then an argument similar to that in the previous paragraph shows that γ_1 is an isomorphism. Suppose $\text{Im}(\theta_0\partial_1^H) \not\subseteq \mathfrak{m}C^{\alpha_1}$. This means the matrix representation of $\theta_0\partial_1^H$ contains a unit; see 1.8. Thus, there exist maps $\rho: C^{\alpha_1} \rightarrow C$ and $\iota: C \rightarrow C^{\alpha_1}$ such that $\rho\theta_0\partial_1^H\iota = \text{id}_C$. This provides a splitting $\partial_1^H\iota$ of $\rho\theta_0: H_0 \rightarrow C$, and so $H_0 \cong C \oplus \ker(\rho\theta_0)$. Finally, the summand $C \oplus 0$ is isomorphic to $\text{Im}(\partial_1^H\iota)$ which is contained in $\text{Im}(\partial_1^H) \subseteq H_0$, contradicting assumption (3).

The fact that γ_0 is an isomorphism now follows as in [4, (8.5)].

Next we show that M admits a resolution satisfying (1)–(3). With the first part of this proof, this will establish part (a). First, note that by Corollary 4.7, there exists a G_C -projective approximation of M

$$Y = 0 \rightarrow K \rightarrow H \rightarrow M \rightarrow 0$$

where H is totally C -reflexive and K admits a bounded $\mathcal{P}_C^f$ -resolution.

If possible, write $H \cong C \oplus H'$ and assume K contains the non-zero C -summand $C \oplus 0$ of H , say $K = C \oplus K'$ for some K' . One checks readily that the compatibility of

the two splittings gives rise to a split exact sequence of complexes, written vertically

$$\begin{array}{ccccccc}
 & 0 & & 0 & & 0 & & 0 \\
 & \downarrow & & \downarrow & & \downarrow & & \downarrow \\
 & Y' & \longrightarrow & K' & \longrightarrow & H' & \longrightarrow & M \longrightarrow 0 \\
 & \downarrow & & \downarrow & & \downarrow & & \downarrow \\
 & Y & \longrightarrow & K & \longrightarrow & H & \longrightarrow & M \longrightarrow 0 \\
 & \downarrow & & \downarrow & & \downarrow & & \downarrow \\
 & X & \longrightarrow & C & \longrightarrow & C & \longrightarrow & 0 \\
 & \downarrow & & \downarrow & & \downarrow & & \downarrow \\
 & 0 & & 0 & & 0 & & 0.
 \end{array}$$

Since Y' is a G_C^f -projective approximation of M , one can repeat this process. Finitely many iterations yield a G_C -approximation $Y_0 = 0 \rightarrow K_0 \rightarrow H_0 \rightarrow M \rightarrow 0$ where K_0 does not contain a nonzero C -summand of H_0 . Lemma 4.9 implies that K admits a minimal $\mathcal{P}_C^f$ -resolution Z . Splicing together Y_0 and Z at K provides a resolution of M satisfying (1)–(3).

Finally, let G be a resolution of M satisfying conditions (1)–(3), and let H be a minimal proper G_C -projective resolution of M . It follows from [11, (1.8)] that G and H are homotopy equivalent. Since G and H are minimal, it follows from Definition 1.9 that they are isomorphic, and so H has the prescribed form. $\square$

ACKNOWLEDGMENTS

I would like to thank my advisor, Sean Sather-Wagstaff, for many stimulating conversations related to this work and for his detailed suggestions. I would also like to thank Lars Winther Christensen, Henrik Holm, Peter Jørgensen, and Greg Piepmeyer and for their comments and suggestions. Finally, I would like to thank Jan Strooker for helpful comments on the exposition.

REFERENCES

1. T. Araya, R. Takahashi, and Y. Yoshino, *Homological invariants associated to semi-dualizing bimodules*, J. Math. Kyoto Univ. **45** (2005), no. 2, 287–306. MR 2161693
2. M. Auslander and M. Bridger, *Stable module theory*, Memoirs of the American Mathematical Society, No. 94, American Mathematical Society, Providence, R.I., 1969. MR 42 #4580
3. M. Auslander and R.-O. Buchweitz, *The homological theory of maximal Cohen-Macaulay approximations*, Mém. Soc. Math. France (N.S.) (1989), no. 38, 5–37, Colloque en l'honneur de Pierre Samuel (Orsay, 1987). MR 1044344 (91h:13010)
4. L. L. Avramov and A. Martsinkovsky, *Absolute, relative, and Tate cohomology of modules of finite Gorenstein dimension*, Proc. London Math. Soc. (3) **85** (2002), 393–440. MR 2003g:16009
5. L. W. Christensen, *Gorenstein dimensions*, Lecture Notes in Mathematics, vol. 1747, Springer-Verlag, Berlin, 2000. MR 2002e:13032
6. ———, *Semi-dualizing complexes and their Auslander categories*, Trans. Amer. Math. Soc. **353** (2001), no. 5, 1839–1883. MR 2002a:13017
7. E. E. Enochs and O. M. G. Jenda, *Gorenstein injective and projective modules*, Math. Z. **220** (1995), no. 4, 611–633. MR 1363858 (97c:16011)
8. E. E. Enochs and O. M. G. Jenda, *Relative homological algebra*, de Gruyter Expositions in Mathematics, vol. 30, Walter de Gruyter & Co., Berlin, 2000. MR 1753146 (2001h:16013)

9. H.-B. Foxby, *Gorenstein modules and related modules*, Math. Scand. **31** (1972), 267–284 (1973). MR 48 #6094
10. E. S. Golod, *G -dimension and generalized perfect ideals*, Trudy Mat. Inst. Steklov. **165** (1984), 62–66, Algebraic geometry and its applications. MR 85m:13011
11. H. Holm, *Gorenstein homological dimensions*, J. Pure Appl. Algebra **189** (2004), no. 1, 167–193. MR 2038564 (2004k:16013)
12. H. Holm and D. White, *Auslander and Bass classes over associative rings*, J. Math. Kyoto Univ. **47** (2007), no. 4, 781808.
13. H. Holm and P. Jørgensen, *Semi-dualizing modules and related Gorenstein homological dimensions*, J. Pure Appl. Algebra **205** (2006), no. 2, 423–445. MR 2203625
14. J. J. Rotman, *An introduction to homological algebra*, Pure and Applied Mathematics, vol. 85, Academic Press Inc. [Harcourt Brace Jovanovich Publishers], New York, 1979. MR 538169 (80k:18001)
15. T. Sharif, S. Sather-Wagstaff and D. White, *Comparison of relative cohomology theories with respect to semidualizing modules*, preprint (2007) [arXiv:math.AC/0706.3635](#).
16. ———, *Gorenstein cohomology in abelian categories*, J. Math Kyoto University, to appear, available from [arXiv:math.AC/0706.3739](#).
17. ———, *Stability of Gorenstein categories*, J. Lond. Math. Soc., to appear, available from [arXiv:math.AC/0703644v2](#)
18. R. Takahashi and D. White, *Homological aspects of semidualizing modules*, J. Commutative Algebra, to appear, available from [arXiv:math.AC/0703643](#).
19. W. V. Vasconcelos, *Divisor theory in module categories*, North-Holland Publishing Co., Amsterdam, 1974, North-Holland Mathematics Studies, No. 14, Notas de Matemática No. 53. [Notes on Mathematics, No. 53]. MR MR0498530 (58 #16637)

DEPARTMENT OF MATHEMATICS, LeCONTE COLLEGE, UNIVERSITY OF SOUTH CAROLINA, 1523 GREENE ST., COLUMBIA, SC, 29208 USA

E-mail address: dwhite@math.sc.edu